




**Catalog Number: JX10076**
**Package size: 25, 100µl**
**Store at: -20°C**

## E-cadherin antibody

<b>Product Name</b>	<b>E-cadherin antibody</b>
<b>Product Number</b>	<b>JX10076</b>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Application</b>	WB, ICC/IF, IHC-P
<b>Species Reactivity</b>	Human
<b>Isotype</b>	IgG
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Storage Buffer</b>	100mM Tris Glycine, 20% Glycerol (pH7) contains 0.025% ProClin 300
<b>Form</b>	Liquid
<b>Recommended Applications Dilutions</b>	Western Blot 1:500-1:1000 Immunocytochemistry / Immunofluorescence 1:200-1:500 Immunohistochemistry (Paraffin) 1:100-1:200
<b>Notes</b>	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

<p>IHC-P analysis of mouse colon tissue section using JX10076 E-cadherin antibody (Green). Dilution: 1:200</p>	<p>IHC-P analysis of human cancer tissue section using JX10076 E-cadherin antibody. Dilution: 1:100</p>	<p>IHC-P analysis of human liver carcinoma tissue section using JX10076 E-cadherin antibody. Dilution: 1:100</p>
		

WB analysis of various sample extracts using JX10076 E-cadherin antibody.  
Loading amount: 60 µg per lane  
Dilution: 1:500

WB analysis of various sample extracts using JX10076 E-cadherin antibody.  
Loading amount: 60 µg per lane  
Dilution: 1:500

WB analysis of various sample extracts using JX10076 E-cadherin antibody.  
Loading amount: 60 µg per lane  
Dilution: 1:500

