


Catalog Number: JX10120

MMP14 / MT1-MMP antibody

Package size: 25, 100µl Store at: -20°C

Product Name	MMP14 / MT1-MMP antibody
Product Number	JX10120
Host	Rabbit
Clonality	Polyclonal
Application	WB, ICC/IF, IHC-P
Species Reactivity	Human, Mouse, Rat
lsotype	IgG
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-
	term storage (1-2 weeks), store at 4° C. For long-term storage, aliquot and store at -
	20°C or below. Avoid multiple freeze-thaw cycles.
Storage Buffer	100mM Tris Glycine, 20% Glycerol (pH7) contains 0.025% ProClin 300
Form	Liquid
Recommended	Western Blot 1:500
Applications Dilutions	Immunocytochemistry / Immunofluorescence 1:150-1:300
	Immunohistochemistry (Paraffin) 1:100-1:200
Notes	Gently mix before use. Optimal concentrations and conditions for each application should
	be determined by the user.

ICC/IF analysis of 4% paraformaldehyde	ICC/IF analysis of 4% paraformaldehyde	WB analysis of various sample extracts
fixed cells using JX10120 MMP14 / MT1-	fixed cells using JX10120 MMP14 / MT1-	using JX10120 MMP14 / MT1-MMP
MMP antibody (Red) counterstained with	MMP antibody (Red) counterstained with	antibody.
DAPI (blue).	DAPI (blue).	Loading amount: 30 µg per lane
Permeabilization: 0.1% NP-40 for 10 min at	Permeabilization: 0.1% NP-40 for 10 min at	Dilution: 1:500
RT	RT	
Dilution: 1:150	Dilution: 1:150	



WB analysis of various sample extracts using JX10120 MMP14 / MT1-MMP antibody.	IHC-P analysis of human cancer tissue section using JX10120 MMP14 / MT1-MMP antibody. Dilution: 1:100	IHC-P analysis of human cancer tissue section using JX10120 MMP14 / MT1-MMP antibody. Dilution: 1:100
Loading amount: 30 µg per lane Dilution: 1:500		
A549 72	Testis carcinoma Liver carcinoma	Breast carcinoma

IHC-P analysis of human breast cancer

tissue section using JX10120 MMP14 /

MT1-MMP antibody.

Dilution: 1:100

Breast carcinoma

