

Catalog Number: JX10135

Package size: 25, 100µl

Store at: -20°C


NFE2L2 / NRF2 antibody

Product Name	NFE2L2 / NRF2 antibody
Product Number	JX10135
Host	Rabbit
Clonality	Polyclonal
Application	WB, ICC/IF, IHC-P
Species Reactivity	Human, Mouse, Rat
Isotype	IgG
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Storage Buffer	100mM Tris Glycine, 20% Glycerol (pH7) contains 0.025% ProClin 300
Form	Liquid
Recommended Applications Dilutions	Western Blot 1:500-1:1000 Immunocytochemistry / Immunofluorescence 1:100-1:200 Immunohistochemistry (Paraffin) 1:100-1:200
Notes	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

WB analysis of various sample extracts using JX10135 NFE2L2 / NRF2 antibody.

Loading amount: 30 µg per lane


Dilution: 1:800



ICC/IF analysis of 4% paraformaldehyde fixed cells using JX10135 NFE2L2 / NRF2 antibody (Red) counterstained with DAPI (blue).

Permeabilization: 0.1% NP-40 for 10 min at RT


Dilution: 1:200












ICC/IF analysis of 4% paraformaldehyde fixed cells using JX10135 NFE2L2 / NRF2 antibody (Red) counterstained with DAPI (blue).

Permeabilization: 0.1% NP-40 for 10 min at RT

Dilution: 1:200




<p>IHC-P analysis of mouse kidney tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>	<p>IHC-P analysis of mouse liver tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:200</p>	<p>IHC-P analysis of mouse lung tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:200</p>
		
<p>IHC-P analysis of mouse tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>	<p>IHC-P analysis of mouse tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>	<p>IHC-P analysis of mouse tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>
		
<p>IHC-P analysis of mouse tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>	<p>IHC-P analysis of mouse tissue section using JX10135 NFE2L2 / NRF2 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>	<p>IHC-P analysis of human cancer tissue section using JX10135 NFE2L2 / NRF2 antibody. Dilution: 1:150</p>
		<p>Breast carcinoma</p> 



IHC-P analysis of human cancer tissue section using JX10135 NFE2L2 / NRF2 antibody.

Dilution: 1:150

Kidney carcinoma



IHC-P analysis of human cancer tissue section using JX10135 NFE2L2 / NRF2 antibody.

Dilution: 1:150

Liver carcinoma

