

Catalog Number: JX10168


SQSTM1 / P62 antibody
Package size: 25, 100µl
Store at: -20°C

Product Name	SQSTM1 / P62 antibody
Product Number	JX10168
Host	Rabbit
Clonality	Polyclonal
Application	WB, ICC/IF, IHC-P
Species Reactivity	Human, Mouse, Rat
Isotype	IgG
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Storage Buffer	100mM Tris Glycine, 1% BSA, 20% Glycerol (pH7) contains 0.025% ProClin 300
Form	Liquid
Recommended Applications Dilutions	Western Blot 1:1000-1:2000 Immunocytochemistry / Immunofluorescence 1:100-1:400 Immunohistochemistry (Paraffin) 1: 100-1:200
Notes	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

WB analysis of various sample extracts using JX10168 SQSTM1 / P62 antibody.

Loading amount: 35 µg per lane


Dilution: 1:1000



ICC/IF analysis of 4% paraformaldehyde fixed cells using JX10168 SQSTM1 / P62 antibody (Red) counterstained with DAPI (blue).

Permeabilization: 0.1% NP-40 for 10 min at RT


Dilution: 1:200










ICC/IF analysis of 4% paraformaldehyde fixed cells using JX10168 SQSTM1 / P62 antibody (Red) counterstained with DAPI (blue).

Permeabilization: 0.1% NP-40 for 10 min at RT

Dilution: 1:100



<p>ICC/IF analysis of 4% paraformaldehyde fixed cells using JX10168 SQSTM1 / P62 antibody (Red) counterstained with DAPI (blue). Permeabilization: 0.1% NP-40 for 10 min at RT Dilution: 1:100</p>	<p>ICC/IF analysis of 4% paraformaldehyde fixed cells using JX10168 SQSTM1 / P62 antibody (Red) counterstained with DAPI (blue). Permeabilization: 0.1% NP-40 for 10 min at RT Dilution: 1:100</p>	<p>IHC-P analysis of mouse tissue section using JX10168 SQSTM1 / P62 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>
 <p style="text-align: center;">SQSTM1 / p62</p> <p style="writing-mode: vertical-rl; transform: rotate(180deg);">HUVEC</p>	 <p style="text-align: center;">SQSTM1 / p62</p> <p style="writing-mode: vertical-rl; transform: rotate(180deg);">MDA-MB-231</p>	
<p>IHC-P analysis of mouse kidney tissue section using JX10168 SQSTM1 / P62 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:200</p>	<p>IHC-P analysis of mouse spleen tissue section using JX10168 SQSTM1 / P62 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:200</p>	<p>IHC-P analysis of mouse tissue section using JX10168 SQSTM1 / P62 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>
		
<p>IHC-P analysis of mouse tissue section using JX10168 SQSTM1 / P62 antibody. Antigen retrieval: Citrate Buffer, pH 6.0 Dilution: 1:100</p>	<p>IHC-P analysis of human colorectal cancer tissue section using JX10168 SQSTM1 / P62 antibody. Dilution: 1:200</p>	
	<p style="text-align: center;">Colorectal carcinoma</p> 