



Catalog Number: JX10189

Package size: 25, 100µl

Store at: -20°C


## VCAM-1 / CD106 antibody

<b>Product Name</b>	VCAM-1 / CD106 antibody
<b>Product Number</b>	JX10189
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Application</b>	WB, ICC/IF, IHC-P
<b>Species Reactivity</b>	Human, Mouse
<b>Isotype</b>	IgG
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Storage Buffer</b>	100mM Tris Glycine, 20% Glycerol (pH7) contains 0.025% ProClin 300
<b>Form</b>	Liquid
<b>Recommended Applications Dilutions</b>	Western Blot 1:500–1:1000 Immunocytochemistry / Immunofluorescence 1:200–1:500 Immunohistochemistry (Paraffin) 1:100–1:200
<b>Notes</b>	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

ICC/IF analysis of 4% paraformaldehyde fixed cells using JX10189 VCAM-1 / CD106 antibody (Red) counterstained with DAPI (blue).


Permeabilization: 0.1% NP-40 for 10 min at RT

Dilution: 1:200




IHC-P analysis of human placenta tissue section using JX10189 VCAM-1 / CD106 antibody (Red).

Dilution: 1:100




IHC-P analysis of human breast cancer tissue section using JX10189 VCAM-1 / CD106 antibody.


Dilution: 1:100




IHC-P analysis of human colorectal cancer tissue section using JX10189 VCAM-1 / CD106 antibody.  
Dilution: 1:100



WB analysis of various sample extracts using JX10189 VCAM-1 / CD106 antibody.  
Loading amount: 50 µg per lane  
Dilution: 1:1000



WB analysis of various sample extracts using JX10189 VCAM-1 / CD106 antibody.  
Loading amount: 50 µg per lane  
Dilution: 1:1000



WB analysis of various sample extracts using JX10189 VCAM-1 / CD106 antibody.  
Loading amount: 50 µg per lane  
Dilution: 1:1000

